



## Length-weight relationships for 11 fish species from the Gulf of Tunis (SW Mediterranean Sea, Tunisia)

MOURAD CHERIF<sup>1</sup>, RAFIK ZARRAD<sup>1</sup>, HOUCINE GHARBI<sup>1</sup>, HECHMI  
MISSAOUI<sup>2</sup> & OTHMAN JARBOUI<sup>1</sup>

<sup>1</sup>National Institute of Sea Sciences and Technologies (I.N.S.T.M), Fishing port 2060, La Goulette, Tunisia. Tel / Fax: (216) 71 735 848. E-mail : mourad.cherif@instm.rnrt.tn

<sup>2</sup>National Agronomic institute of Tunisia (I.N.A.T)

**Abstract.** In this study, length–weight relationships of 11 commercial fish species from the Gulf of Tunis: *Mullus barbatus*, *Scomber scombrus*, *Sparus aurata*, *Boops boops*, *Spicara maena*, *Diplodus annularis*, *Merluccius merluccius*, *Trachurus trachurus*, *Trachurus mediterraneus*, *Pagellus erythrinus* and *Lithognathus mormyrus* were presented. The values of the slope *b* in the length - weight relationship ranged from 2.674 to 3.368 and intercepts between 0.0021 and 0.0515.

**Key words:** Fishery resources, fish morphometry, ichthyofauna of the Mediterranean Sea.

**Resumo. Relações comprimento-peso para 11 espécies de peixes do Golfo de Tunes (Mar Mediterraneo Sudocidental, Tunisia).** No presente estudo, as relações comprimento-peso para 11 espécies comerciais de peixes do Golfo de Tunis: *Mullus barbatus*, *Scomber scombrus*, *Sparus aurata*, *Boops boops*, *Spicara maena*, *Diplodus annularis*, *Merluccius merluccius*, *Trachurus trachurus*, *Trachurus mediterraneus*, *Pagellus erythrinus* e *Lithognathus mormyrus* foram apresentadas. O valor do coeficiente angular *b* nas relações comprimento – peso variou entre 2.674 e 3.368 e o valor de interseção da reta no eixo *Y* variou entre 0.0021 e 0.0515.

**Palavras-chave:** Recursos pesqueiros, morfometria de peixes, ictiofauna do mar mediterrânea.


The Gulf of Tunis provides a significant proportion of the overall marine fish production in Tunisia and is considered one of the most important fishery grounds of the Tunisian coast (Azouz 1973, Gharbi 1980, Zarrad *et al.*, 2001). The Gulf of Tunis is also known as an important spawning and nursery ground for several fish species (Hattour 1991, Zarrad *et al.*, 2003).

Length and weight data are a useful and standard result of fish sampling programs. These data are needed to estimate growth rates, length and age structures, and other components of fish population dynamics (Kolher *et al.*, 1995). Length-weight relationships allow fisheries scientists to convert growth-in-length equations to growth-in-weight in stock assessment models (Dulčić & Kraljević 1996, Gonçalves *et al.*, 1997, Morato *et al.*, 2001, Stergiou & Moutopoulos 2001, Özaydin *et al.*, 2007), estimate biomass from length frequency distributions (Anderson & Gutreuter 1983, Petrakis

& Stergiou 1995, Dulčić & Kraljević 1996), and calculate fish condition (Petrakis & Stergiou 1995). Length-weight relationships are also useful for comparing life history and morphological aspects of populations inhabiting different regions (Gonçalves *et al.*, 1997, Stergiou & Moutopoulos 2001). Despite the utility of length-weight relationship in fisheries science and the importance of the Gulf of Tunis for Tunisian fisheries, information about the length-weight relationships of fish species in Tunis Gulf is scarce and incomplete.

The present study presents estimates of the length weight relationships for 11 species of the families Carangidae, Mullidae, Scombridae, Sparidae and Merlucciidae.

Data were collected during twelve monthly trawl surveys conducted in February 2003 to January 2004 in the gulf of Tunis (Fig 1). Twelve hauls were operated monthly, based on a random sampling method, covering an area of 570 km<sup>2</sup>



**Figure 1.** Map of Gulf of Tunis

between 40 and 100 m of depth, on professional boats. These experimental surveys were operated monthly on professional boats, covering an area of 570 km<sup>2</sup> between 40 and 100 m of depth. The vessel used was equipped with a Tunisian shrimp's trawl (modified Gulf of Mexico trawl) with 52 mm stretched mesh in the wing and 40 mm in the cod-end. Each haul lasted 120 min at an average speed of 3 knots.

In the laboratory specimens were sorted by sex, measured to the nearest 1 mm (total length, TL) and weighed to the nearest 0.1 g (weight, W). The relationship between the length and weight of a fish is usually expressed by the equation  $W=aL^b$  (Ricker 1973) where W is body weight (g), L is total length (cm), *a* is the intercept and *b* is the slope (fish growth rate) (Beverton & Holt 1996). The parameters *a* and *b* of the length-weight relationships were estimated by the least-square method based on the predictive or Type I linear

regression model (Sokal & Rohlf 1981), using W as the dependent variable and L as the independent variable, the data analysis has given that for each specie, the number of outliers is far less than 10% of total data and thus the least square model is satisfactory for each set of data (Chen & Jackson 2000). The degree of adjustment of the model studied was assessed by the correlation coefficient (*r*). Student's t-test was applied to verify whether the declivity of regression (constant "b") presented a significant difference of 3.0, indicating the type of growth: isometric (*b*=3.0), positive allometric (*b*>3.0) or negative allometric (*b*<3.0) (Spiegel 1991). In all cases a statistic significance of 5% was adopted.

A total of 5533 individual fish length and weight observations were recorded for the 11 fish species analyzed in this study. The estimated parameters and length characteristics of the length-weight relationship are given in Table I.

**Table I.** Descriptive statistics and estimated parameters of the length-weight relationship for 11 fish species from the Gulf of Tunis. min, minimum; max, maximum; S.E., standard error; n, the sample size; a, the intercept of the relationship; b, the slope of the relationship; r, coefficient of correlation.

Species	Sex	n	Length characteristics (cm)					Parameters					t-test
			mean	SE	min	max	a	b	SE (b)	r			
<i>Mullus barbatus</i> (Linnaeus, 1758)	males	664	12,8	0,12	6,3	21	0,004	3,28	0,0314	0,99	4,237*		
	females	1099	16,4	0,07	9,7	25	0,007	3,12	0,0521	0,97	5,073*		
<i>Sparus aurata</i> (Linnaeus, 1758)	both	1763	14,9	0,07	6,3	25	0,005	3,23	0,0072	0,99	3,291*		
	males	124	19	0,1	16	22	0,052	2,8	0,0651	0,92	3,096*		
<i>Diplodus annularis</i> (Linnaeus, 1758)	females	271	17	0,06	17	23	0,014	2,72	0,0398	0,93	3,566*		
	both	395	19,2	0,05	16	23	0,03	2,67	0,0152	0,91	4,381*		
<i>Pagellus erythrinus</i> (Linnaeus, 1758)	males	183	13,2	0,09	9,6	17	0,033	2,72	0,0557	0,92	3,022*		
	females	287	13,1	0,07	10	17	0,01	2,86	0,0723	0,92	2,873*		
<i>Spicara maena</i> (Linnaeus, 1758)	both	470	13	0,52	9,6	17	0,012	2,9	0,0752	0,93	2,895*		
	males	285	19,1	0,16	11	29	0,01	2,84	0,0358	0,97	2,012*		
<i>Merluccius merluccius</i> (Linnaeus, 1758)	females	613	16,8	0,08	11	25	0,018	2,84	0,0637	0,97	2,395*		
	both	898	17,2	0,84	7,4	29	0,017	2,85	0,0239	0,98	2,208*		
<i>Boops boops</i> (Linnaeus, 1758)	males	121	16	0,17	9,9	19	0,03	2,82	0,0448	0,97	1,708 <sup>ns</sup>		
	females	209	14,9	0,11	11	19	0,012	2,94	0,0333	0,96	1,695 <sup>ns</sup>		
<i>Trachurus trachurus</i> (Linnaeus, 1758)	both	330	15,4	0,09	9,5	20	0,014	2,87	0,0328	0,96	1,493 <sup>ns</sup>		
	males	187	22,4	0,35	16	43	0,004	3,17	0,0354	0,98	3,511*		
<i>Trachurus mediterraneus</i> (Steindachner, 1863)	females	340	26,1	0,24	17	41	0,005	3,11	0,0284	0,98	2,996*		
	both	527	24,8	0,21	16	43	0,004	3,13	0,0232	0,98	4,147*		
<i>Lithognathus mormyrus</i> (Linnaeus, 1758)	males	71	18,2	0,24	15	23	0,003	3,01	0,0827	0,97	0,245 <sup>ns</sup>		
	females	172	19,5	0,19	12	26	0,009	2,98	0,0458	0,97	0,232 <sup>ns</sup>		
<i>Scomber scombrus</i> (Linnaeus, 1758)	both	243	18,5	0,14	12	26	0,007	3,06	0,0278	0,98	0,255 <sup>ns</sup>		
	males	85	19,3	0,32	13	32	0,007	3,00	0,0242	0,98	0,553 <sup>ns</sup>		
<i>Trachurus mediterraneus</i> (Steindachner, 1863)	females	144	19,1	0,17	15	25	0,007	2,98	0,0168	0,97	0,287 <sup>ns</sup>		
	both	229	17,5	0,17	9	32	0,007	2,98	0,0279	0,99	0,749 <sup>ns</sup>		
<i>Lithognathus mormyrus</i> (Linnaeus, 1758)	males	182	18,9	0,28	8,8	30	0,007	2,99	0,0206	0,98	0,201 <sup>ns</sup>		
	females	280	19,6	0,18	12	28	0,009	2,9	0,0135	0,96	0,231 <sup>ns</sup>		
<i>Scomber scombrus</i> (Linnaeus, 1758)	both	462	17,9	0,2	8,8	30	0,007	3,01	0,0236	0,99	0,166 <sup>ns</sup>		
	males	28	18,5	0,28	15	21	0,005	2,92	0,0457	0,96	0,854 <sup>ns</sup>		
<i>Scomber scombrus</i> (Linnaeus, 1758)	females	89	18,8	0,15	15	22	0,01	2,98	0,0398	0,98	0,778 <sup>ns</sup>		
	both	117	18,7	0,13	15	22	0,009	2,84	0,0291	0,98	0,852 <sup>ns</sup>		
<i>Scomber scombrus</i> (Linnaeus, 1758)	males	44	24,4	0,49	17	32	0,002	3,35	0,0385	0,98	3,177*		
	females	55	26,1	0,31	18	32	0,006	3,18	0,0429	0,99	2,386*		
both	99	25,2	0,21	17	32	0,002	3,37	0,1658	0,99	3,557*			

\*: p&lt;0.05; ns: p&gt;0.05

The sample size ranged from 99 individuals for *Scomber scombrus* to 1763 for *Mullus barbatus*. The  $r$  values ranged from 0.82 for *Sparus aurata* to 0.99 for *Trachurus trachurus*, and all regressions were highly significant. Values of  $b$  ranged from 2.674 for *Sparus aurata* to 3.368 for *Scomber scombrus*. A high degree of positive correlation between total length and total weight of all 11 fish species is indicated by high values of correlation coefficient  $r$ . The estimated values of  $b$  were close to 3.0 ( $p > 0.05$ ) showing isometric growth for *Lithognathus mormyrus*, *Boops boops*, *Spicara maena*, *Trachurus trachurus* and *Trachurus mediterraneus*.

The values of the exponent  $b$  for males, females and combined sexes were significantly ( $p < 0.05$ ) higher than 3.0 exhibited a positive allometric growth for *Mullus barbatus*, *Merluccius merluccius*, and *Scomber scombrus*. The slopes for male, female and both sexes for *Sparus aurata*, *Diplodus annularis* and *Pagellus erythrinus* showed a significant difference ( $p < 0.05$ ) lower than 3.0 reflecting a negative allometric growth.

Even though the change of  $b$  values depends primarily on the shape and fatness of the species, various factors may be responsible for the differences in parameters of the length/weight relationships among seasons and years, such as temperature, salinity, food (quantity, quality and size), sex, time of year and stage of maturity (Pauly 1984, Sparre 1992). According to Bagenal & Tesch (1978), Gonçalves *et al.*, (1997), Taskavak & Bilecenoglu (2001) and Özeydin *et al.*, (2007), the parameter  $b$ , unlike the parameter  $a$ , may vary seasonally, and even daily, and between habitats. Thus, the length-weight relationship in fish is affected by a number of factors including gonad maturity, sex, diet, stomach fullness, health, and preservation techniques as well as season and habitat, none of which were taken into consideration in the present study.

The information gained in the present survey can enable fish biologists to derive weight estimates for the Gulf of Tunis fishes that are measured but not weighed. The length-weight parameters hereby reported may be of considerable use in ongoing studies of catches in Tunisian commercial fisheries.

### Acknowledgments

We are grateful to Zahara Youssef and Ajengui Amara from the National Institute of Sea Sciences and Technologies for their help in collecting samples.

### References

- Azouz, A. 1973. Les fonds chalutables de la région Nord de la Tunisie. **Bulletin de l'Institut National Scientifique et Technique d'Océanographie et de Pêche de Salammbô**, 2(4): 473-563.
- Bagenal, T. B. & Tesch, F. W. 1978. **Age and growth**. In: T. Bagenal (Editor). **Methods for Assessment of Fish Production in Fresh Waters**. IBP Handbook No. 3, 3rd ed. Blackwell Science Publications. pp. 101-136.
- Beverton, R. J. H. & Holt, S. J. 1996. **On the Dynamics of Exploited Fish Populations**. Chapman and Hall, London.
- Chen, Y. & Jackson, D. A. 2000. An empirical study on estimators for linear regression analyses in fisheries and ecology. **Fisheries Research**, 49: 193-206.
- Dulčić, J. & Kraljević, M. 1996. Weight-length relationships for 40 fish species in the eastern Adriatic (Croatian waters). **Fisheries Research**, 28: 243-251.
- Gharbi, H. 1980. Contribution à l'étude biologique et dynamique des rougets (*Mullus barbatus Linnaeus, 1758* et *Mullus surmuletus Linnaeus, 1758*) des côtes tunisiennes. **Thèse doctorat de spécialité (3<sup>ème</sup> cycle de biologie marine)** Université de Tunis, Fac. Sic. Tunis: 100 p.
- Gonçalves, J. M. S., Bentes, L., Lino, P. G., Ribeiro, J., Canario, A. V. M. & Erzini, K. 1997. Weight-length relationships for selected fish species of the small-scale demersal fisheries of the south and south-west coast of Portugal. **Fisheries Research**, 30: 253-256.
- Hattour, A. 1991. Le chalutage dans les eaux tunisiennes : réalités et considérations législatives dans les golfe de Tunis et de Gabès. **Notes. Bulletin de l'Institut National Scientifique et Technique d'Océanographie et de Pêche de Salammbô**, 1 : 26 p.
- Kolher, N., Casey, J. & Turner, P. 1995. Length-weight relationships for 13 species of sharks from the western North Atlantic. **Fishery Bulletin**, 93: 412-418.
- Morato, T. P., Afonso, P., Lourinho, P., Barreiros, J. P., Santos, R. S. & Nash, R. D. M. 2001. Length-weight relationships for 21 coastal fish species of the Azores, north-eastern Atlantic. **Fisheries Research**, 50: 297-302.
- Özeydin, O. & Taskavak, E. 2007. Length-weight relationships for 47 fish species from Izmir Bay (eastern Aegean Sea, Turkey). **Acta Adriatica**, 47(2): 211-216.

- Pauly, D. 1984. Fish population dynamics in tropical waters: A manual for use with programmable calculators. **ICLARM Studies and Reviews 8**. ICLARM, Manila, Philippines. 325 pp.
- Petrakis, G. & Stergiou, K. I. 1995. Weight length relationships for 33 fish species in Greek waters. **Fisheries Research**, 21: 465-469.
- Ricker, W. E. 1973. Linear regressions in fishery research. **Journal of the Fisheries Research Board of Canada**, 30: 409-434.
- Sokal, R. R. & Rohlf, F. G. 1981. **Biometry**. W.H. Freeman & Co, 843 p.
- Sparre, P. 1992. Introduction to Tropical Fish Stock Assessment. **Part I- Manual. FAO Fisheries Technical Paper 306/1**. Rev 1. 1992. Rome.
- Spiegel, M. R. 1991. **Théorie et applications de la statistique**.: McGraw-Hill, Paris, 358p.
- Stergiou, K. I. & Moutopoulos, D. K. 2001. A review of length-weight relationships of fishes from Greek marine waters. **Naga, ICLARM Quart**, 24(1-2): 23-39.
- Taskavak, E. & Bilecenoglu, M. 2001. Length weight relationships for 18 Lessepsian (Red Sea) immigrant fish species from the eastern Mediterranean coast of Turkey. **Journal of the Marine Biological Association of the United Kingdom**, 81: 895-896.
- Zarrad, R., Gharbi, H. & Missaoui, H. 2001. Détermination de l'effort optimal de chalutage benthique dans le golfe de Tunis. **Bulletin de l'Institut National des Sciences et Technologies de la Mer de Salammbô**, 28: 3-7
- Zarrad, R., El Abed, A., M'rabet, R., Missaoui, H. & Romdhane, M. S. 2003. Distribution spatiale de l'ichtyoplancton en été et en automne et conditions environnementales dans le golfe de Tunis. **Bulletin de l'Institut National des Sciences et Technologies de la Mer de Salammbô**, 30:39-47.

Received August 2007

Accepted November 2007

Published online January 2008